



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Mobile technology offers novel insights on control and treatment of allergic rhinitis. The MASK study

**Citation for published version:**

MASK study group 2019, 'Mobile technology offers novel insights on control and treatment of allergic rhinitis. The MASK study', *Journal of Allergy and Clinical Immunology*.  
<https://doi.org/10.1016/j.jaci.2019.01.053>

**Digital Object Identifier (DOI):**

[10.1016/j.jaci.2019.01.053](https://doi.org/10.1016/j.jaci.2019.01.053)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Journal of Allergy and Clinical Immunology

**Publisher Rights Statement:**

This is the author's peer-reviewed manuscript as accepted for publication.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.


**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Accepted Manuscript

Mobile technology offers novel insights on control and treatment of allergic rhinitis.  
The MASK study



Annabelle Bédard, MD, Xavier Basagaña, PhD, Josep M. Anto, PhD, Judith Garcia-Aymerich, MD, Philippe Devillier, MD, Sylvie Arnavielhe, PhD, Anna Bedbrook, BSc, Gabrielle L. Onorato, MSc, Wienczyslawa Czarlewski, MD, Ruth Murray, PhD, Rute Almeida, PhD, Joao Fonseca, MD, Elisio Costa, PhD, Joao Malva, MD, Mario Morais-Almeida, MD, Ana Maria Pereira, MD, Ana Todo-Bom, MD, Enrica Menditto, PhD, Cristiana Stellato, MD, Maria Teresa Ventura, MD, Alvaro A. Cruz, MD, Rafaël Stelmach, MD, Jane da Silva, MD, Désirée Larenas-Linnemann, MD, José M. Fuentes-Pérez, MD, Yunuen R. Huerta-Villalobos, MD, Regina Emuzyte, MD, Violeta Kvedariene, MD, Arunas Valiulis, MD, Piotr Kuna, MD, Boleslaw Samolinski, MD, Ludger Klimek, MD, Ralph Mösges, MD, Oliver Pfaar, MD, Sara Shamai, MD, Isabelle Annesi-Maesano, MD, Isabelle Bosse, MD, Pascal Demoly, MD, Jean-François Fontaine, MD, Vicky Cardona, MD, Joaquim Mullol, MD, Antonio Valero, MD, Regina E. Roller-Wirnsberger, MD, Peter Valentin Tomazic, MD, Niels H. Chavannes, MD, Wytske J. Fokkens, MD, Sietze Reitsma, MD, Mike Bewick, MD, Dermot Ryan, MD, Aziz Sheikh, MD, Tari Haahtela, MD, Sanna Toppila-Salmi, MD, Erkka Valovirta, MD, Michael Makris, MD, Nikos G. Papadopoulos, MD, Emmanuel P. Prokopoulos, MD, Fotis Psarros, MD, Cemal Cingi, MD, Bilun Gemicioğlu, MD, Arzu Yorgancioglu, MD, Sinthia Bosnic-Anticevich, PhD, Robyn E. O'Hehir, MD, Claus Bachert, MD, Peter W. Hellings, MD, Benoit Pugin, PhD, Carsten Bindeslev-Jensen, MD, Esben Eller, MD, Ingrid Kull, PhD, Erik Melén, MD, Magnus Wickman, MD, Gert De Vries, MSc, Michiel van Eerd, MSc, Ioana Agache, MD, Ignacio J. Ansotegui, MD, Mark S. Dykewicz, MD, Thomas Casale, MD, Dana Wallace, MD, Susan Wasserman, MD, Daniel Laune, PhD, Jean Bousquet, MD, and the MASK study group

PII: S0091-6749(19)30426-9

DOI: <https://doi.org/10.1016/j.jaci.2019.01.053>

Reference: YMAI 13953

To appear in: *Journal of Allergy and Clinical Immunology*

Received Date: 12 November 2018

Revised Date: 5 January 2019

Accepted Date: 23 January 2019

Please cite this article as: Bédard A, Basagaña X, Anto JM, Garcia-Aymerich J, Devillier P, Arnavielhe S, Bedbrook A, Onorato GL, Czarlewski W, Murray R, Almeida R, Fonseca J, Costa E, Malva J, Morais-Almeida M, Pereira AM, Todo-Bom A, Menditto E, Stellato C, Ventura MT, Cruz AA, Stelmach R, da Silva J, Larenas-Linnemann D, Fuentes-Pérez JM, Huerta-Villalobos YR, Emuzyte R, Kvedariene V, Valiulis A, Kuna P, Samolinski B, Klimek L, Mösges R, Pfaar O, Shamaï S, Annesi-Maesano I, Bosse I, Demoly P, Fontaine J-F, Cardona V, Mullol J, Valero A, Roller-Wirnsberger RE, Tomazic PV, Chavannes NH, Fokkens WJ, Reitsma S, Bewick M, Ryan D, Sheikh A, Haahtela T, Toppila-Salmi S, Valovirta E, Makris M, Papadopoulos NG, Prokopakis EP, Psarros F, Cingi C, Gemicioğlu B, Yorgancioglu A, Bosnic-Anticevich S, O'Hehir RE, Bachert C, Hellings PW, Pugin B, Bindeslev-Jensen C, Eller E, Kull I, Melén E, Wickman M, De Vries G, van Eerd M, Agache I, Ansotegui IJ, Dykewicz MS, Casale T, Wallace D, Wasserman S, Laune D, Bousquet J, and the MASK study group, Mobile technology offers novel insights on control and treatment of allergic rhinitis. The MASK study, *Journal of Allergy and Clinical Immunology* (2019), doi: <https://doi.org/10.1016/j.jaci.2019.01.053>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Mobile technology offers novel insights on control and treatment of allergic rhinitis. The MASK study

Annabelle Bédard, MD, (1-3), Xavier Basagaña, PhD, (1-3), Josep M Anto, PhD, (1-3), Judith Garcia-Aymerich, MD, (1-3), Philippe Devillier, MD, (5), Sylvie Arnavielhe, PhD, (6), Anna Bedbrook, BSc, (7), Gabrielle L Onorato, MSc, (7), Wienczyslawa Czarlewski, MD, (8), Ruth Murray, PhD, (9), Rute Almeida, PhD, (10), Joao Fonseca, MD, (10), Elisio Costa, PhD, (11), Joao Malva, MD, (12), Mario Morais-Almeida, MD, (13), Ana Maria Pereira, MD, (14), Ana Todo-Bom, MD, (15), Enrica Menditto, PhD, (16), Cristiana Stellato, MD, (17), Maria Teresa Ventura, MD, (18), Alvaro A Cruz, MD, (19), Rafaël Stelmach, MD, (20), Jane da Silva, MD, (21), Désirée Larenas-Linnemann, MD, (22), José M Fuentes-Pérez, MD (23), Yunuen R Huerta-Villalobos, MD, (23), Regina Emuzyte, MD, (24), Violeta Kvedariene, MD, (25), Arunas Valiulis, MD, (26), Piotr Kuna, MD, (27), Boleslaw Samolinski, MD, (28), Ludger Klimek, MD, (29), Ralph Mösges, MD, (30), Oliver Pfaar, MD, (32), Sara Shamai, MD, (30), Isabelle Annesi-Maesano, MD, (33), Isabelle Bosse, MD, (34), Pascal Demoly, MD, (35), Jean-François Fontaine, MD, (36), Vicky Cardona, MD, (37), Joaquim Mullol, MD, (38), Antonio Valero, MD, (39), Regina E Roller-Wirnsberger, MD, (40), Peter Valentin Tomazic, MD, (41), Niels H Chavannes, MD, (42), Wytke J Fokkens, MD, (43), Sietze Reitsma, MD, (43), Mike Bewick, MD, (44), Dermot Ryan, MD, (45), Aziz Sheikh, MD, (46), Tari Haahtela, MD, (47), Sanna Toppila-Salmi, MD, (47), Erkkä Valovirta, MD, (48), Michael Makris, MD, (49), Nikos G Papadopoulos, MD, (50), Emmanuel P Prokopakis, MD, (51), Fotis Psarros, MD, (52), Cemal Cingi, MD, (53), Bilun Gemicioğlu, MD, (54), Arzu Yorgancioglu, MD, (55), Sinthia Bosnic-Anticevich, PhD, (56), Robyn E O'Hehir, MD, (57), Claus Bachert, MD, (58), Peter W Hellings, MD, (59), Benoit Pugin, PhD (60), Carsten Bindslev-Jensen, MD, (61), Esben Eller, MD, (61), Ingrid Kull, PhD, (62,63), Erik Melén, MD, (63), Magnus Wickman, MD, (64), Gert De Vries, MSc, (65), Michiel van Eerd, MSc, (65), Ioana Agache, MD, (66), Ignacio J Ansotegui, MD, (67), Mark S Dykewicz, MD, (68), Thomas Casale, MD, (69), Dana Wallace, MD, (70), Susan Wasserman, MD, (71), Daniel Laune, PhD, (6), Jean Bousquet, MD, (7, 31, 72), and the MASK study group.

1. ISGlobal, Barcelona, Spain.
2. Universitat Pompeu Fabra (UPF), Barcelona, Spain.
3. CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain.
4. Universitat Pompeu Fabra (UPF), Barcelona, Spain.
5. UPRES EA220, Pôle des Maladies des Voies Respiratoires, Hôpital Foch, Université Paris-Saclay, Suresnes, France.
6. KYomed INNOV, Montpellier, France.
7. MACVIA-France, Fondation partenariale FMC VIA-LR, Montpellier, France.
8. Medical Consulting Czarlewski, Levallois, France.
9. Director, Medical Communications Consultant, MedScript Ltd, Dundalk, Co Louth, Ireland and New Zealand, Honorary Research Fellow, OPC, Cambridge, UK.
10. Center for Health Technology and Services Research - CINTESIS, Faculdade de Medicina, Universidade do Porto; and Medida, Lda Porto, Portugal.
11. UCIBIO, REQUINTE, Faculty of Pharmacy and Competence Center on Active and Healthy Ageing of University of Porto (Porto4Ageing), Porto, Portugal.
12. Institute of Biomedical Imaging and Life Sciences (IBILI), Faculty of Medicine, University of Coimbra, Portugal; Ageing@Coimbra EIP-AHA Reference Site, Portugal.
13. Allergy Center, CUF Descobertas Hospital, Lisbon, Portugal
14. Allergy Unit, CUF-Porto Hospital and Institute; Center for Research in Health Technologies and information systems CINTESIS, Universidade do Porto, Portugal.
15. Imunoalergologia, Centro Hospitalar Universitário de Coimbra and Faculty of Medicine, University of Coimbra, Portugal.

16. CIRFF, Federico II University, Naples, Italy.
17. Department of Medicine, Surgery and Dentistry "Scuola Medica Salernitana", University of Salerno, Salerno, Italy.
18. University of Bari Medical School, Unit of Geriatric Immunoallergology, Bari, Italy.
19. ProAR – Nucleo de Excelencia em Asma, Federal University of Bahia, Brasil and WHO GARD Planning Group, Brazil.
20. Pulmonary Division, Heart Institute (InCor), Hospital da Clinicas da Faculdade de Medicina da Universidade de Sao Paulo, Sao Paulo, Brazil.
21. Department of Internal Medicine and Allergy Clinic of Professor Polydoro Ernani de São Thiago University Hospital, Federal University of Santa Catarina (UFSC), Florianopolis-SC, Brazil.
22. Center of Excellence in Asthma and Allergy, Hospital Médica Sur, México City, Mexico.
23. Hospital General Region 1, Dr Carlos Mc Gregor Sanchez Navarro" IMSS. Mexico City, Mexico
24. Clinic of Children's Diseases, Faculty of Medicine, Vilnius University, Vilnius, Lithuania.
25. Faculty of Medicine, Vilnius University, Vilnius, Lithuania.
26. Vilnius University Institute of Clinical Medicine, Clinic of Children's Diseases, and Institute of Health Sciences, Department of Public Health, Vilnius, Lithuania; European Academy of Paediatrics (EAP/UEMS-SP), Brussels, Belgium.
27. Division of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Poland.
28. Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Poland.
29. Center for Rhinology and Allergology, Wiesbaden, Department of Otorhinolaryngology, Head and Neck Surgery, Universitätsmedizin Mannheim, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany.
30. Institute of Medical Statistics, and Computational Biology, Medical Faculty, University of Cologne, Germany and CRI-Clinical Research International-Ltd, Hamburg, Germany.
31. University Hospital, Montpellier, France
32. Department of Otorhinolaryngology, Head and Neck Surgery, Section for Rhinology and Allergy, University Hospital Marburg, Philipps-Universität, Marburg, Germany.
33. Epidemiology of Allergic and Respiratory Diseases, Department Institute Pierre Louis of Epidemiology and Public Health, INSERM and UPMC Sorbonne Université, Medical School Saint Antoine, Paris, France
34. Allergist, La Rochelle, France.
35. Department of Respiratory Diseases, Montpellier University Hospital, France
36. Allergist, Reims, France.
37. Allergy Section, Department of Internal Medicine, Hospital Vall 'dHebron & ARADyAL research network, Barcelona, Spain.
38. Rhinology Unit & Smell Clinic, ENT Department, Hospital Clínic; Clinical & Experimental Respiratory Immunoallergy, IDIBAPS, CIBERES, University of Barcelona, Spain.
39. Pneumology and Allergy Department CIBERES and Clinical & Experimental Respiratory Immunoallergy, IDIBAPS, University of Barcelona, Spain.
40. Medical University of Graz, Department of Internal Medicine, Graz, Austria.
41. Department of ENT, Medical University of Graz, Austria
42. Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, The Netherlands
43. Department of Otorhinolaryngology, Amsterdam University Medical Centres, AMC, Amsterdam, the Netherlands.
44. iQ4U Consultants Ltd, London, UK.
45. Honorary Clinical Research Fellow, Allergy and Respiratory Research Group, The University of Edinburgh, Edinburgh, UK
46. The Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK.
47. Skin and Allergy Hospital, Helsinki University Hospital, Helsinki, Finland.
48. Department of Lung Diseases and Clinical Immunology, University of Turku and Terveystalo allergy clinic, Turku, Finland.
49. Allergy Unit "D Kalogeromitros", 2nd Dpt of Dermatology and Venereology, National & Kapodistrian University of Athens, "Attikon" University Hospital, Greece.
50. Center for Pediatrics and Child Health, Institute of Human Development, Royal Manchester Children's Hospital, University of Manchester, Manchester M13 9WL, UK Allergy Department, 2nd

- Pediatric Clinic, Athens General Children's Hospital "P&A Kyriakou," University of Athens, Athens 11527, Greece
51. Department of Otorhinolaryngology University of Crete School of Medicine, Heraklion, Greece.
  52. Allergy Department, Athens Naval Hospital, Athens, Greece.
  53. Eskisehir Osmangazi University, Medical Faculty, ENT Department, Eskisehir, Turkey.
  54. Department of Pulmonary Diseases, Istanbul University-Cerrahpasa, Cerrahpasa Faculty of Medicine, Istanbul, Turkey.
  55. Department of Pulmonary Diseases, Celal Bayar University, Faculty of Medicine, Manisa, Turkey and GARD Executive Committee.
  56. Woolcock Institute of Medical Research, University of Sydney and Woolcock Emphysema Centre and Local Health District, Glebe, NSW, Australia.
  57. Department of Allergy, Immunology and Respiratory Medicine, Alfred Hospital and Central Clinical School, Monash University, Melbourne, Victoria, Australia; Department of Immunology, Monash University, Melbourne, Victoria, Australia.
  58. Upper Airways Research Laboratory, ENT Dept, Ghent University Hospital, Ghent, Belgium.
  59. Dept of Otorhinolaryngology, Univ Hospitals Leuven, Belgium, and Academic Medical Center, Univ of Amsterdam, The Netherlands and Euforea, Brussels, Belgium.
  60. European Forum for Research and Education in Allergy and Airway Diseases (EUFOREA), Brussels, Belgium.
  61. Department of Dermatology and Allergy Centre, Odense University Hospital, Odense Research Center for Anaphylaxis (ORCA), Odense, Denmark.
  62. Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet, and Sachs' Children and Youth Hospital, Södersjukhuset, Stockholm, Sweden
  63. Sachs' Children and Youth Hospital, Södersjukhuset, Stockholm and Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden.
  64. Centre for Clinical Research Sörmland, Uppsala University, Eskilstuna, Sweden.
  65. Peercode BV, Geldermalsen, The Netherlands.
  66. Transylvania University Brasov, Brasov, Romania.
  67. Department of Allergy and Immunology, Hospital Quirón Bizkaia, Erandio, Spain.
  68. Section of Allergy and Immunology, Saint Louis University School of Medicine, Saint Louis, Missouri, USA.
  69. Division of Allergy/Immunology, University of South Florida, Tampa, Fla.
  70. Nova Southeastern University, Fort Lauderdale, Florida, USA.
  71. Department of Medicine, Clinical Immunology and Allergy, McMaster University, Hamilton, Ontario, Canada.
  72. INSERM U 1168, VIMA : Ageing and chronic diseases Epidemiological and public health approaches, Villejuif, Université Versailles St-Quentin-en-Yvelines, UMR-S 1168, Montigny le Bretonneux, France and Euforea, Brussels, Belgium

**Short title: Treatment of allergic rhinitis using an App**

## **Author for correspondence**

Professor Jean Bousquet  
 CHU Arnaud de Villeneuve, 371 Avenue du Doyen Gaston Giraud, 34295 Montpellier Cedex 5,  
 France Tel +33 611 42 88 47, Fax :+33 467 41 67 01 [jean.bousquet@orange.fr](mailto:jean.bousquet@orange.fr)



## Conflict of interest

Dr. Devillier reports personal fees from Sanofi-Aventis, GlaxoSmithKline, Astra Zeneca, Chiesi, Meda Pharma, Menarini, outside the submitted work.

Dr. Almeida reports grants from Project NORTE-01-0145-FEDER-000016 (NanoSTIMA) by North Portugal Regional Operational Programme (NORTE 2020), under the Portugal 2020 Partnership Agreement, and through the European Regional Development Fund (ERDF), during the conduct of the study.

Dr. Todo-Bom reports grants and personal fees from GSK, Mundipharma Novartis, personal fees from Teva pharma, personal fees from Astrazeneca, grants from Leti, Bial, outside the submitted work.

Dr. Cruz reports grants and personal fees from Astrazeneca, grants from GSK, personal fees from Boehringer Ingelheim, CHIESI, NOVARTIS, Eurofarma, MEDA Pharma, Boston Scientific, outside the submitted work.

Dr. Larenas Linnemann reports personal fees from Amstrong, Astrazeneca, Boehringer Ingelheim, Chiesi, DBV Technologies, Grunenthal, GSK, MEDA, Menarini, MSD, Novartis, Pfizer, Sanofi, Siegfried, UCB, grants from Sanofi, Astrazeneca, Novartis, UCB, GSK, TEVA, Boehringer Ingelheim, Chiesi, outside the submitted work.

Dr. Stelmach reports grants from São Paulo Research Foundation, MSD, grants and personal fees from Novartis, grants, personal fees and non-financial support from AstraZeneca, Chiesi, personal fees and non-financial support from Boehringer Ingelheim, outside the submitted work.

Dr. Larenas Linnemann reports personal fees from Amstrong, Astrazeneca, Boehringer Ingelheim, Chiesi, DBV Technologies, Grunenthal, GSK, MEDA, Menarini, MSD, Novartis, Pfizer, Sanofi, Siegfried, UCB, grants from Sanofi, Astrazeneca, Novartis, UCB, GSK, TEVA, Boehringer Ingelheim, Chiesi, outside the submitted work.

Dr. Kvedariene has received payment for consultancy from GSK and for lectures from StallergensGreer, Berlin-Chemie outside the submitted work.

Dr. Kuna reports personal fees from Adamed, Boehringer Ingelheim, AstraZeneca, Chiesi, FAES, Berlin Chemie, Novartis, Polpharma, Allergopharma, outside the submitted work.

Dr. Ralph Mösges reports personal fees from ALK, allergopharma, Allergy Therapeutics, Hexal, personal fees from Servier, personal fees from Klosterfrau, Stada, UCB, Friulchem, grants from ASIT biotech, Nuvo, Bayer, FAES, GSK, MSD, Johnson&Johnson, Meda, Optima, Ursapharm, BitopAG, Hulka, grants and personal fees from Bencard, grants from Leti, Stallergenes, grants, personal fees and non-financial support from Lofarma, non-financial support from Roxall, from Atmos, from Bionorica, Otonomy, Ferrero, personal fees and non-financial support from Novartis, outside the submitted work.

Dr. Pfaar reports grants and personal fees from ALK-Abelló, Allergopharma, Stallergenes Greer, HAL Allergy Holding B.V./HAL Allergie GmbH, Bencard Allergie GmbH/Allergy Therapeutics, Lofarma, grants from Biomay, Nuvo, Circassia, Glaxo Smith Kline, personal fees from Novartis Pharma, MEDA Pharma, Indoor Biotechnologies, Pohl-Boskamp, outside the submitted work.

Dr. Haahtela reports personal fees from Mundipharma, Novartis, and Orion Pharma, outside the submitted work.

Dr. Toppila-Salmi reports other from Biomedical Systems Ltd., Roche Ltd., grants from Erkkö Foundation, outside the submitted work.

Dr. Papadopoulos reports grants from Gerolymatos, personal fees from Hal Allergy B.V., Novartis Pharma AG, Menarini, Hal Allergy B.V., Mylan, outside the submitted work.

Dr. Bosnic-Anticevich reports personal fees from Teva, Boehringer Ingelheim, Sanofi, GSK, AstraZeneca, outside the submitted work.

Bachert reports personal fees from Meda, Stallergenes and ALK (speaker).

Dr. Ansotegui reports personal fees from Hikma, Roxall, Astra Zeneca, Menarini, UCB, Faes Farma, Sanofi, Mundipharma, outside the submitted work.

J Bousquet reports personal fees and other from Chiesi, Cipla, Hikma, Menarini, Mundipharma, Mylan, Novartis, Sanofi-Aventis, Takeda, Teva, Uriach, outside the submitted work. other from Kyomed.

Dr. Wallace reports other from Mylan Pharmaceutical Company, outside the submitted work; and I am co-chair of the AAAAI/ACAAI Joint Task Force on Practice Parameters. I helped to develop the Rhinitis GRADE document published 12/2017.

The rest of the authors declare that they have no relevant conflicts of interest

ACCEPTED MANUSCRIPT



## Abstract

**Background:** Mobile health may be used to generate innovative insights into optimizing treatment to improve allergic rhinitis control.

**Objectives:** A cross-sectional real world observational study was undertaken in 22 countries to complement a pilot study and bring novel information on medication use, disease control and work productivity in everyday life of patients with allergic rhinitis.

**Methods:** A mobile phone app (*Allergy Diary*, freely available Google Play and Apple stores) was used to collect data of daily visual analogue scales (VAS) for (i) overall allergic symptoms, (ii) nasal, ocular and asthma symptoms, (iii) work, as well as (iv) medication use using a treatment scroll list including all allergy medications (prescribed and over-the-counter (OTC)) customized for 22 countries. The four most common intra-nasal medications containing intra-nasal corticosteroids and eight oral H1-antihistamines were studied.

**Results:** 9,122 users filled in 112,054 days of VAS in 2016 and 2017. The assessment of days was informative. The control of days with rhinitis differed between no [best control], single [good control for intranasal corticosteroid-treated days] or multiple treatments [worst control]. Users with the worst control increased the range of treatments being used. The same trend was found for asthma, eye symptoms and work productivity. Differences between oral H1-antihistamines were found.

**Conclusions:** This study confirms the usefulness of the *Allergy Diary* in accessing and assessing patient behavior in allergic rhinitis. This observational study using a very simple assessment tool (VAS) on a mobile phone had the potential to answer questions previously thought infeasible.

## Capsule summary

Most rhinitis patients use on-demand treatment when they are not controlled. Control was worse with increasing medications. Real life data may not be aligned with guidelines.

## Clinical implications

A behavioural disconnection was found in the study since patients are not adherent to treatment and treat themselves on-demand when they are not controlled whereas the vast majority of physicians prescribe long-term treatment to achieve control. Shared-decision making is essential.

## Key words

Allergic rhinitis, anti-histamines, asthma, conjunctivitis, corticosteroids, mobile health, MASK, treatment

## Abbreviations

AR: Allergic rhinitis  
 AzeFlu : Intranasal azelastine-fluticasone propionate  
 CET : Cetirizine  
 DL: Desloratadine  
 FEXO: Fexofenadine  
 FF: Fluticasone Furoate  
 FP: Fluticasone Propionate  
 INCS: Intranasal corticosteroid  
 INN: International Nonproprietary Names  
 LEVOCET: Levocetirizine  
 Lora: Loratadine  
 MASK-rhinitis (Mobile Airways Sentinel Network for allergic rhinitis)  
 Mometasone Furoate (MF)  
 OAH: Oral H<sup>1</sup>-anti-histamine  
 RCT: Randomized controlled trial  
 visual analogue scales (VAS)

## Introduction

The treatment of allergic rhinitis (AR) is complex as many drugs are available in oral and/or topical formulations. Many guidelines for AR are evidence-based and have led to a better management of AR. However, guidelines are mostly based on randomized controlled trials (RCTs), typically undertaken on highly selected populations, often with limited/unclear generalizability to routine care contexts (1, 2). They propose to increase treatment to achieve disease control (i.e. sleep, social and school/work impairment) that is the ultimate aim of the treatment. Intra-nasal corticosteroids represent the most effective AR treatment for most patients, but their effect is relatively slow, taking several hours (3) and many patients prefer oral medications. A formulation of fluticasone propionate (FP) and azelastine (AzeFlu) is more effective than INCS alone (4) and has the advantage of acting within minutes (5). Patients are poorly adherent to treatment and often self-medicate (6, 7). They want more effective and fast acting treatments. Observational real-life studies are therefore needed to complement RCTs in order to better understand the efficacy of INCS-containing medications since they do not select patients and report their behavior.

MASK-rhinitis (Mobile Airways Sentinel NetworK for allergic rhinitis), an information and communications technology (ICT) system centered around the patient (8-12) operational in 23 countries, uses a treatment scroll list including all medications customized for each country and a visual analogue scales (VAS) to assess rhinitis control. A pilot study in over 2,900 users allowed differentiation between treatments (13). Patients did not necessarily use treatment on a daily basis in a regular way but appeared to increase treatment use when their symptom's control worsens. However, the pilot study needs to be confirmed with a larger number of users and more medications tested.

The present cross-sectional observational study was undertaken in 9,122 users in 22 countries (data collection was just started in Argentina) to confirm the pilot study (13) using the same methods and to bring novel information on medication use, and associated disease control, work productivity (14) and allergic multimorbidity (13). The study was focused firstly on the four most commonly used intra-nasal medications containing intra-nasal corticosteroids: Fluticasone Furoate (FF), Fluticasone Propionate (FP), Mometasone Furoate (MF) and AZeFlu. We did not perform the same analysis with oral H1-antihistamines as they are often associated with INCS and many patients would have been analysed twice. In the second analysis, we examined some widely used oral H1-antihistamines: Bilastine, Cetirizine (CET), Desloratadine (DL), Ebastine, Fexofenadine (FEXO), Levocetirizine (LEVOCET), Loratadine (Lora) and Rupatadine. In the first analysis, we compared days with single treatment with days with multiple treatments. In the second analysis, we just used days with a single treatment.

## Methods

### Users

All consecutive users from January 1, 2016 to December 31, 2017 were included with no exclusion criteria according to methods previously described (13, 14).

### Setting

Users from 22 countries filled in the *Allergy Diary* (Table 1). Data collection was just started in Argentina and not included

### Ethics

The Allergy Diary is CE1. CE marking is a certification mark that indicates conformity with health, safety, and environmental protection standards for products made in the EU and meets the essential requirements of all relevant European Medical Device Directives (15). CE1 includes sterile and non-sterile products and assess whether the device has a measuring function.

The data were anonymized including data related to geolocalization using k-anonymity (16).

An independent Review Board approval was not required since the study is observational and users agreed to have their data analysed (terms of use).

### Allergy Diary

Geolocalized users assess their daily symptom control using the touchscreen functionality on their smart phone to click on five consecutive VAS scores (i.e. general, nasal and ocular symptoms, asthma and work). Users input their daily medications using a scroll list which contains all country-specific OTC and prescribed medications available for each country (Figure 1 online). The list has been populated using IMS data.

Days reported by users included days with or without treatment.

The present study is another *Allergy Diary* study. Some of the raw data used in the first paper (up to November 2016) (13) were used in this study, but analyses differed.

### Selection of medications

The International Nonproprietary Names (INN) classification was used for drug nomenclature

(17). Monotherapy was defined as days when only one single medication for rhinitis was reported. AzeFlu contains two drugs but, as it is a fixed combination it was considered as monotherapy. Co-medication was defined as days with two or more medications for rhinitis. Asthma medications were not considered in co-medication.

## Size of the study

In this study, all registered users were included to obtain the best possible estimates for the specified time window. From the pilot study, numbers tested largely exceed those needed to find significant differences in the full set analysis (13). However, we did not consider medications with a sample size under 1,000 days of reporting.

## Statistical methods

A non-Gaussian distribution was found for the data. Non-parametric tests and medians (and percentiles) were used. Correction for multiple testing was made when appropriate.

Some users reported VAS scores more than once a day. In the pilot study, we found that the highest reported value should be used and we followed this study (13). We however tested in an exploratory analysis VAS levels in duplicates and multiplicates.

## Analysis of the data

We conducted, as previously published (13), separate analyses using the full-set of data and data on just the first day of reporting.

In the first analysis, only users who reported no treatment or treatment by the intra-nasal FF, FP, MF and AZeFlu were studied (Figure 2 online). Those receiving other INCS were excluded. For co-medication, we initially selected second generation oral H1-antihistamines (OAH): CET, DL, Ebastine, FEXO, LEVOCET, LORA and Rupatadine (Group + OAH). There are many other OAH, but we did not consider them since their pharmacologic properties vary widely and they were not often used. We considered two other groups in INCS users for co-medication: users who reported OAH and another medication (Group OAH + other) and users who reported another medication (+ Other). Users who reported other medications but no INCS were not analyzed. As a primary end point, using the full data set, we studied median VAS global measured ("Overall how much are your allergic symptoms bothering you today?") levels for days with FF, FP, MF and AZeFlu and for days without medications. The primary and secondary end points were analyzed using the Kruskal-Wallis test and Wilcoxon and Mann-Whitney test with Dunn-Bonferroni's post hoc analysis to correct for multiple testing.

Moreover, we analysed the data using three cutoffs: VAS <20/100 (controlled days), VAS 20-49 (days with moderate control), VAS  $\geq$ 50 (days with poor control) according to a consensus (18) and available data of the pilot study (13, 14). The same analyses were conducted for the first day of VAS report. Secondary end-points included VAS eye, asthma and work.

In the second analysis, we compared days with monotherapy for the most common OAH: CET, DL, Ebastine, FEXO, LEVOCET, LORA and Rupatadine monotherapy. We did not consider other OAH with a sample size under 1,000 days (or close to this number). We only compared VAS global measured. The mean number of days of reporting was considered for each treatment.

We then performed exploratory analyses to investigate whether there are temporal patterns in the reporting of VAS in the app users. We assessed the VAS levels on: (i) days with more than 1 VAS reported, (ii) the first day of reporting and first day of new reporting in users with non-consecutive data, (iii) days without treatment followed by a day with treatment and (iv) days with treatment followed by a day without treatment.

## Results

### Demographic characteristics

The study included 9,122 users. Roughly 5% of users did not report their age and were ascribed to “zero”. Users ranged in age from zero to 92 years (mean, SD:  $32.4 \pm 15.2$  years). There were 54.7% women and 45.3% men. The age repartition is given in Figure 3 online.

A total of 112,054 days was recorded. Duplicates or multiplies for the same day were found in 14,767 days. Global VAS was not recorded in 754 (0.8 %) days with App data reported. There were 52,706 (54.6%) days without treatment and 18,117 days with the targeted INCS (Figure 1).

### Analysis of VAS global measured

On visual inspection, no clear trajectory of VAS could be easily identified, as users reported erratically their VAS and treatment data. Figure 4 online reports trajectories for French users as an example.

In the figure each user is identified by a member identifier number (vertical axis) and each user's trajectory is represented horizontally by dots - each dot representing a day of VAS recording).

Results are reported in Table 2, Figures 2 and 3.

### **Analysis of VAS global measured on days without treatment and days with INCS treatment**

The first day of reporting, VAS levels were reported by 4,991 users without treatment, 1,395 users with OAH and 1,281 users with INCS treatment (Table 2). The percentage of users with single treatment ranged from 34.0% (FP), 39.2% (MF), 40.5% (FF) and 59.6% (AzeFlu). Days with INCS alone had similar median VAS levels (35 to 44).

For the full data set of 96,533 days, VAS levels were reported by 6,236 users without treatment, 3,664 users with OAH and 2,575 users with INCS treatment (Table 2). Monotherapy was reported 45 to 55% of the days (FF or MF versus AzeFlu – Figure 2). For monotherapy, median VAS levels ranged from 5 (FF) to 23.5 (FP). For day 1 and the full data set, the same trend was found in INCS treated users: lowest median levels were found for monotherapy, increased levels with co-medication by OAH and highest levels for co-medication with OAH + other treatments (Figure 3). Variable levels of VAS were observed for co-medication with other treatments. The numbers of days of co-medication with another INCS are too low to make any comparison (Table 2).

### **Analysis of VAS global measured on days with OAH treatment alone**

The first day of reporting, days with no treatment or those with INCS in monotherapy had similar median VAS levels (34 to 44). On the other hand, there were some variations for OAH in monotherapy. LEVOCET days had a median VAS level intermediate between untreated or INCS-treated days and the other OAH. For the full data set of 96,533 days, median VAS levels of days with INCS were lower than those of days with OAH but Bilastine, FEXO, LEVOCET and Rupatadine had levels similar to those of INCS (Table 2).

Apart from days with FP treatment (low numbers), the mean numbers of days of reporting medications per user ranged from 4.00 (CET) to 8.98 (AzeFlu).

### **Analyses of VAS for eye, asthma and work**

Analyses of VAS eye, asthma and work are reported in Figures 5A, B and C online supplement. Trends for the three secondary end points are similar to those of VAS global measured, i.e. low median levels similar to untreated days for the single treatment, increased levels with co-medication by OAH and highest levels for co-medication with OAH + other medication, and the highest percentage of users with single treatment observed for AzeFlu. Fewer users reported VAS work, but the trends were similar.



## Exploratory analyses investigating potential temporal patterns in the reporting of VAS

### Assessment of duplicates or multiplicates for day 1

Days with 2 or more VAS levels reported at least 1 hour apart within the same day were selected. The dataset included 1,576 days for VAS global measured. A significantly higher VAS was found at second reporting compared to the first. When the data were stratified by the type of treatment recorded at first entry (no treatment, AzeFlu FF, MF and FP), these findings were only significant for days with no treatment. No difference was found for days with (any) treatment (Table 1 online).

### VAS levels depending on consecutive and non-consecutive data

There were 4,132 users with at least two non-consecutive calendar days of VAS reported (n=89,473 days in total). The global VAS levels measured on day 1 were found to be significantly higher when compared to the global VAS levels measured on the first day of new reporting (i.e. or first non-consecutive calendar day reported), regardless of the presence/type of treatment (Table 3).

The distribution of global VAS on the 391 consecutive couple of calendar days consisting of a day without treatment followed by a day with treatment showed a non-significant increased level in treated days (median [p25-75] =23 [11-49] to 28 [14-50], (p=0.07, Wilcoxon W test).

The distribution of global VAS on the 350 consecutive couple of calendar days consisting of a day with treatment followed by a day without treatment showed a significant decreased level in untreated days (median [p25-75] =23 [13-45] to 20 [9-38], (p=0.01 Wilcoxon W test).

## Discussion

A pilot study using a very simple assessment (VAS) on a cell phone in 2,871 users who filled in 17,091 days suggested that an App may give novel information concerning the treatment of AR (13). However, the sample size was possibly too small to draw definite conclusions. This study in a larger sample (9,111 users in 22 countries, 97,287 days) confirms the findings of the pilot study showing that, in real life, the assessment of days can inform on patient's treatment and bring novel insight on the behaviour of AR patients towards treatment and novel concepts for change management of AR (19). The control of days differs between no treatment (best control), single treatment or co-medication (worst control). This study showed for the first time that the same trends were observed for global symptoms, ocular symptoms, asthma and work productivity. This study suggests contrary behaviour between physicians and patients since the

range of treatments was increased in those with poor control whereas, according to guidelines, physicians are recommended to increase the treatment to achieve control. This major gap in AR treatment may explain the overall low level of satisfaction of severe AR patients reported in many studies.

## Strengths and limitations

The current study has many strengths including larger numbers, multiple countries, range of treatments studied and patient/person-generated data.

As for all studies using participatory data, potential biases include (i) the likelihood of sampling bias likely present, difficult to assess generalizability of the study, (ii) outcome misclassification that cannot be assessed and, by definition due to ethical problems, there very little information on patient (or day) characteristics. App users are not representative of all patients with rhinitis. The issue of potential selection bias was limited by the fact that we considered days and not patients in the analyses.

As in other studies (13, 20), we used days in a cross-sectional analysis because there is no clear pattern of treatment and a longitudinal study was not feasible since users mostly use the App intermittently. Although this observation may differ from RCTs, our study is a real-life approach.

For this study, other biases should be considered. The diagnosis of AR was not supported by a physician but was a response to the question: “Do you have allergic rhinitis? Yes/No”. There may therefore be some users with non-allergic rhinitis who may have responded “Yes” to the question. There are potential measurement biases when using apps including collection of information, education of the patient, availability and ability to use a smartphone (13). Users self-identified themselves as having AR without confirmation of the diagnosis. Precise patient characterization is impossible using an App, but every observational study using the *Allergy Diary* was able to identify days with poor control or criteria of severity (20-24). Adherence to treatment is impossible to prove as users do not report data all days and users may not report all medications used. Nonetheless, mobile technology is becoming an important tool to better understand and manage AR and brings novel information that were not available with other methods (20-26).

Asthma was assessed using a single VAS largely validated in rhinitis (27). In asthma, VAS was shown to be an effective measure of control (28). In the present study, we did not investigate specific symptoms or perform any pulmonary function test. Thus, it is possible that some users

may have misunderstood the question or overestimated the disease. However, the results are extremely consistent.

We only considered days and not patients' trajectories because these are highly variable, patients using auto-medication depending on AR control as previously shown (13).

Longitudinal capture is very challenging with this App but this appears to be the case for all Apps. Patient's engagement with digital health in real world scenarios is usually lower than in RCTs. Although this is a limitation in relation to causal inference, it suggests that a new methodological approach is needed. It appears that treatment trajectories are specific for almost each user and most users have gaps in their treatment when they are well controlled.

### **Interpretation of the results and generalizability**

This real world assessment of the *Allergy Diary* using VAS allows assessment of treatment efficacy by days, which represents real-life estimation of AR control and likely reflects real-life better than patients' assessments at regular intervals since (i) it is known that AR is a highly variable disease, and control varies widely between days in relation to allergen and environmental exposure, (ii) patients are rarely adherent to their treatment, (iii) patients often stop treatment when they feel better and (iv) patients increase their treatment when uncontrolled.

VAS scores were greater on days with treatment than on days without treatment. This study confirms the study of the pilot one (13) in which, median VAS levels on days without treatment were similar in users who never reported any medication use and in those who were occasionally treated. Moreover, in a small sample, it was found that consecutive days under treatment are less well controlled than days without treatment. In INCS-treated users, days with a single treatment were better controlled than days with multiple treatments. An important message from this paper is that, overall, in real life, patients treat themselves when they suffer from symptoms and stop their treatment when they are controlled. This accords with previous data (29, 30). This study, using objective data, confirmed that adherence is poor. Most AR patients may have mild and/or intermittent disease that does not need a regular treatment to achieve control. The concept of pro-active medication and patient participation (31) - the patient starting treatment when experiencing symptoms and continuing for a few days after getting control - may be of great interest and could be tested with the App. In asthma, self-guided treatment was found to be of interest (31-33). Such real-life findings may ultimately affect the way in which guidelines are constructed to align them more with human behaviour. We have already initiated a program entitled Change management in rhinitis and asthma (19) in which

we propose to develop next-generation care pathways and test the recommendations of GRADE guidelines in AR (3, 4) according to real-world evidence using data of MASK. A first meeting was held at the Pasteur Institute, Paris (December 3, 2018) to provide guidance for their development.

This observational study made it possible to differentiate OAH and INCS, confirming known data, (34) and was able to differentiate between OAH. LEVOCET was found to be the most effective OAH confirming clinical experience. On the other hand, CETI appeared not to have been as effective. However, there were a large number of generics for CETI and this could be studied when more users will be available. This study could also differentiate the three medications containing INCS: FF, MF and MP-AZeFlu and confirm previous studies (35)(36) extending our understanding of how AR treatment is used. RCTs showed that MP-AzeFlu is more effective than single components available in pharmacies (37) or components using the same formulation (38).

The same trends for INCS-containing medications were observed for VAS global measured, eye, asthma and work. However, the percentages of well-controlled, controlled and poorly-controlled days differed indicating the independence of data already observed. Moreover, data on work are extremely important to facilitate an economic evaluation of treatments.

An important result is that VAS on day 1 was higher than any other consecutive/non-consecutive day. This indicates that patients start using the App when symptoms are uncontrolled. This is one specificity of analysing app data and should be considered in studies that assess the control of allergic diseases in relation to risk factors such as air pollutants and allergen exposure.

## Conclusions

Real world data (RWD) and real-world evidence (RWE) are playing an increasing role in health care decisions supporting clinical trial designs and observational studies to generate innovative and new treatment approaches. These data hold potential to answer questions previously thought infeasible (39) such as the true patient's attitude towards treatment. This observational study shows highly consistent results between different outcomes (VAS levels) and brings novel concepts for the management of allergic diseases. When the patient experiences increased symptom, indicating a loss of control, he/she increases the number of medications used that day. A total behavioural disconnection was found since most patients treat themselves on-demand when they are not controlled whereas the vast majority of physicians prescribe long-term treatment to achieve control. Shared decision making may offer a more rewarding approach AR

560 management. The results of this paper will be of importance for the implementation of the  
561 MASK Good Practice recently recognized by DG Santé.  
562  
563

## References

1. Price D, Smith P, Hellings P, Papadopoulos N, Fokkens W, Muraro A, et al. Current controversies and challenges in allergic rhinitis management. *Expert Rev Clin Immunol*. 2015;1-13.
2. Travers J, Marsh S, Williams M, Weatherall M, Caldwell B, Shirtcliffe P, et al. External validity of randomised controlled trials in asthma: to whom do the results of the trials apply? *Thorax*. 2017;62(3):219-23.
3. Brozek JL, Bousquet J, Agache I, Agarwal A, Bachert C, Bosnic-Anticevich S, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) Guidelines - 2016 Revision. *J Allergy Clin Immunol*. 2017;140(4):950-8.
4. Dykewicz MS, Wallace DV, Baroody F, Bernstein J, Craig T, Finegold I, et al. Treatment of seasonal allergic rhinitis: An evidence-based focused 2017 guideline update. *Ann Allergy Asthma Immunol*. 2017;119(6):489-511 e41.
5. Bousquet J, Meltzer EO, Couroux P, Koltun A, Kopietz F, Munzel U, et al. Onset of Action of the Fixed Combination Intranasal Azelastine-Fluticasone Propionate in an Allergen Exposure Chamber. *J Allergy Clin Immunol Pract*. 2018;6(5):1726-32.
6. Bosnic-Anticevich S, Kritikos V, Carter V, Yan KY, Armour C, Ryan D, et al. Lack of asthma and rhinitis control in general practitioner-managed patients prescribed fixed-dose combination therapy in Australia. *J Asthma*. 2018;55(6):684-94.
7. Tan R, Cvetkovski B, Kritikos V, Price D, Yan K, Smith P, et al. Identifying the hidden burden of allergic rhinitis (AR) in community pharmacy: a global phenomenon. *Asthma Res Pract*. 2017;3:8.
8. Bourret R, Bousquet J, J M, T C, Bedbrook A, P D, et al. MASK rhinitis, a single tool for integrated care pathways in allergic rhinitis. *World Hosp Health Serv*. 2015;51(3):36-9.
9. Bousquet J, Schunemann HJ, Fonseca J, Samolinski B, Bachert C, Canonica GW, et al. MACVIA-ARIA Sentinel NetworK for allergic rhinitis (MASK-rhinitis): the new generation guideline implementation. *Allergy*. 2015;70(11):1372-92.
10. Bousquet J, Hellings PW, Agache I, Bedbrook A, Bachert C, Bergmann KC, et al. ARIA 2016: Care pathways implementing emerging technologies for predictive medicine in rhinitis and asthma across the life cycle. *Clin Transl Allergy*. 2016;6:47.
11. Bousquet J, Anto JM, Annesi-Maesano I, Dedeu T, Dupas E, Pepin JL, et al. POLLAR: Impact of air POLLution on Asthma and Rhinitis; a European Institute of Innovation and Technology Health (EIT Health) project. *Clin Transl Allergy*. 2018;8:36.
12. Bousquet J, Arnavielhe S, Bedbrook A, Bewick M, Laune D, Mathieu-Dupas E, et al. MASK 2017: ARIA digitally-enabled, integrated, person-centred care for rhinitis and asthma multimorbidity using real-world-evidence. *Clin Transl Allergy*. 2018;8:45.
13. Bousquet J, Arnavielhe S, Bedbrook A, Alexis-Alexandre G, Eerd Mv, Murray R, et al. Treatment of allergic rhinitis using mobile technology with real world data: The MASK observational pilot study. *Allergy*. 2018:sous presse.
14. Bousquet J, Bewick M, Arnavielhe S, Mathieu-Dupas E, Murray R, Bedbrook A, et al. Work productivity in rhinitis using cell phones: The MASK pilot study. *Allergy*. 2017;72(10):1475-84.
15. Council Directive 93/42/EEC of 14 June 1993 concerning medical devices. 1993L0042 — EN — 11.10.2007 — 005.001 — 1. <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CONSLEG:1993L0042:20071011:EN:PDF>. 1993.
16. Samreth D, Arnavielhe S, Ingenrieth F, Bedbrook A, Onorato GL, Murray R, et al. Geolocation with respect to personal privacy for the Allergy Diary app - a MASK study. *World Allergy Organ J*. 2018;11(1):15.
17. Kopp-Kubel S. International Nonproprietary Names (INN) for pharmaceutical substances. *Bull World Health Organ*. 1995;73(3):275-9.
18. Bousquet J, Schunemann HJ, Hellings PW, Arnavielhe S, Bachert C, Bedbrook A, et al. MACVIA clinical decision algorithm in adolescents and adults with allergic rhinitis. *J Allergy Clin Immunol*. 2016;138(2):367-74 e2.



19. Bousquet J, Hellings PW, Agache I, Amat F, Annesi-Maesano I, Ansotegui IJ, et al. ARIA Phase 4 (2018): Change management in allergic rhinitis and asthma multimorbidity using mobile technology. *J Allergy Clin Immunol*. 2018;pii: S0091-6749(18)31359-9. doi: 10.1016/j.jaci.2018.08.049.
20. Caimmi D, Baiz N, Tanno LK, Demoly P, Arnavielhe S, Murray R, et al. Validation of the MASK-rhinitis visual analogue scale on smartphone screens to assess allergic rhinitis control. *Clin Exp Allergy*. 2017;47(12):1526-33.
21. Bousquet J, Caimmi DP, Bedbrook A, Bewick M, Hellings PW, Devillier P, et al. Pilot study of mobile phone technology in allergic rhinitis in European countries: the MASK-rhinitis study. *Allergy*. 2017;72(6):857-65.
22. Bousquet J, Arnavielhe S, Bedbrook A, Fonseca J, Morais Almeida M, Todo Bom A, et al. The Allergic Rhinitis and its Impact on Asthma (ARIA) score of allergic rhinitis using mobile technology correlates with quality of life: The MASK study. *Allergy*. 2018;73(2):505-10.
23. Bousquet J, Devillier P, Anto JM, Bewick M, Haahtela T, Arnavielhe S, et al. Daily allergic multimorbidity in rhinitis using mobile technology: a novel concept of the MASK study. *Allergy*. 2018;73(9):1763-74.
24. Bousquet J, VandenPlas O, Bewick M, Arnavielhe S, Bedbrook A, Murray R, et al. The Work Productivity and Activity Impairment Allergic Specific (WPAI-AS) Questionnaire Using Mobile Technology: The MASK Study. *J Invest Allergol Clin Immunol*. 2018;28(1):42-4.
25. Bonini M. Electronic health (e-Health): emerging role in asthma. *Curr Opin Pulm Med*. 2017;23(1):21-6.
26. Pizzulli A, Perna S, Florack J, Pizzulli A, Giordani P, Tripodi S, et al. The impact of telemonitoring on adherence to nasal corticosteroid treatment in children with seasonal allergic rhinoconjunctivitis. *Clin Exp Allergy*. 2014;44(10):1246-54.
27. Klimek L, Bergmann KC, Biedermann T, Bousquet J, Hellings P, Jung K, et al. Visual analogue scales (VAS): Measuring instruments for the documentation of symptoms and therapy monitoring in cases of allergic rhinitis in everyday health care: Position Paper of the German Society of Allergology (AeDA) and the German Society of Allergy and Clinical Immunology (DGAKI), ENT Section, in collaboration with the working group on Clinical Immunology, Allergology and Environmental Medicine of the German Society of Otorhinolaryngology, Head and Neck Surgery (DGHNOKHC). *Allergo J Int*. 2017;26(1):16-24.
28. Ohta K, Jean Bousquet P, Akiyama K, Adachi M, Ichinose M, Ebisawa M, et al. Visual analog scale as a predictor of GINA-defined asthma control. The SACRA study in Japan. *J Asthma*. 2013;50(5):514-21.
29. Kremer B, Klimek L, Gulicher D, Degen M, Mosges R. Sequential therapy with azelastine in seasonal allergic rhinitis. Deutsche Rhinitis Studiengruppe (German Rhinitis Study Group). *Arzneimittelforschung*. 1999;49(11):912-9.
30. Salo T, Peura S, Salimaki J, Maasilta P, Haahtela T, Kauppi P. Need for medication and stuffy nose predict the severity of allergic rhinitis. *Asia Pac Allergy*. 2016;6(2):133-5.
31. Lahdensuo A, Haahtela T, Herrala J, Kava T, Kiviranta K, Kuusisto P, et al. Randomised comparison of cost effectiveness of guided self management and traditional treatment of asthma in Finland. *BMJ*. 1998;316(7138):1138-9.
32. Lahdensuo A, Haahtela T, Herrala J, Kava T, Kiviranta K, Kuusisto P, et al. Randomised comparison of guided self management and traditional treatment of asthma over one year. *Bmj*. 1996;312(7033):748-52.
33. McDonald VM, Gibson PG. Asthma self-management education. *Chron Respir Dis*. 2006;3(1):29-37.
34. Brozek JL, Bousquet J, Baena-Cagnani CE, Bonini S, Canonica GW, Casale TB, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines: 2010 revision. *J Allergy Clin Immunol*. 2010;126(3):466-76.



- 671 35. Meltzer EO, Wallace D, Dykewicz M, Shneyer L. Minimal Clinically Important Difference  
672 (MCID) in Allergic Rhinitis: Agency for Healthcare Research and Quality or Anchor-  
673 Based Thresholds? *J Allergy Clin Immunol Pract.* 2016;4(4):682-8 e6.
- 674 36. Bachert C, Bousquet J, Hellings P. Rapid onset of action and reduced nasal hyperreactivity:  
675 new targets in allergic rhinitis management. *Clin Transl Allergy.* 2018;8:25.
- 676 37. Hampel FC, Ratner PH, Van Bavel J, Amar NJ, Daftary P, Wheeler W, et al. Double-blind,  
677 placebo-controlled study of azelastine and fluticasone in a single nasal spray delivery  
678 device. *Ann Allergy Asthma Immunol.* 2010;105(2):168-73.
- 679 38. Carr W, Bernstein J, Lieberman P, Meltzer E, Bachert C, Price D, et al. A novel intranasal  
680 therapy of azelastine with fluticasone for the treatment of allergic rhinitis. *J Allergy Clin*  
681 *Immunol.* 2012;129(5):1282-9 e10.
- 682 39. Sherman RE, Anderson SA, Dal Pan GJ, Gray GW, Gross T, Hunter NL, et al. Real-World  
683 Evidence - What Is It and What Can It Tell Us? *N Engl J Med.* 2016;375(23):2293-7.
- 684
- 685

**Table 1. Country and number of users recording Visual Analogue Scale score using the Allergy Diary in the full data set**

Country	VAS measurements (days)				Total
	1	2 to 7	8 to 14	>14	
Austria	226 (56.6%)	121	16	36	399
Australia	49 (49.0%)	30	10	11	100
Belgium	48 (49.5%)	35	5	9	97
Brazil	572 (55.9%)	323	67	62	1024
Canada	6 (35.3%)	7	3	1	17
Czech Republic	1 (20.0%)	0	1	3	5
Denmark	37 (45.1%)	29	4	12	82
Finland	117 (44.8%)	93	25	26	261
France	319 (61.3%)	147	19	35	520
Germany	208 (39.8%)	141	35	139	523
Greece	47 (23.7%)	43	24	84	198
Italy	554 (44.6%)	389	87	213	1243
Lithuania	59 (17.7%)	89	52	134	334
Mexico	101 (13.0%)	207	128	343	779
Netherland	167 (53.9%)	94	23	26	310
Poland	286 (54.9%)	159	28	48	521
Portugal	647 (49.2%)	505	64	100	1316
Spain	129 (30.5%)	124	53	117	423
Sweden	33 (39.3%)	34	6	11	84
Switzerland	247 (64.0%)	111	11	17	386
Turkey	81 (52.6%)	42	10	21	154
UK	148 (42.8%)	104	46	48	346
<b>Total</b>	<b>4082 (44.7%)</b>	<b>2827 (31.0%)</b>	<b>717 (7.9%)</b>	<b>1496 (16.4%)</b>	<b>9122</b>

691 Table 2: Results of VAS global measured

	Day 1		Full set (96,533 days)		
	N days	Median [p25-p75]	N days [users]	Median [p25-p75]	Mean number of days per user
No treatment	4991	34 [10-60]	52706 [6236]	8 [0-26]	8.45
Bilastine*	128	48 [19-69.5]	1563 [261]	16 [6-37]	6.00
Cetirizine*	350	52 [28-70]	2169 [545]	22 [9-50]	4.00
Desloratadine*	300	50 [26-71]	2085 [504]	21 [8-46]	4.14
Ebastine*	115	50 [26-72]	980 [201]	23 [9-48]	4.88
Fexofenadine*	112	55 [32.5-71.5]	1128 [183]	14 [8-35]	6.17
Levocetirizine*	149	43 [16-67]	1512 [260]	14 [5-28]	5.81
Loratadine*	175	49 [28-72]	1680 [344]	21 [10-39]	4.88
Rupatadine*	66	49 [23-63]	1138 [146]	18 [5-36]	7.69
FF	176	35 [19.5-58.5]	2182 [336]	5 [0-27]	6.49
+ OAH	129	51 [22-66]	1317 [247]	21 [4-45]	5.33
+ OAH + other	38	64 [49-77]	307 [80]	48 [24-63]	3.84
+ other (no OAH)	84	53.5 [28-72]	968 [168]	23 [9-47]	5.76
+ other INCS	7	50 [4-90]	113 [16]	61 [26-95]	7.06
AzeFlu	155	37 [16-60]	2722 [303]	13 [3-29]	8.98
+ OAH	49	58 [40-73]	994 [113]	17 [7-40]	8.72
+ OAH + other	12	54 [26-80]	174 [33]	31 [9-60]	5.27
+ other (no OAH)	37	40 [21-65]	871 [98]	22 [11-42]	8.89
+ other INCS	7	50 [33-77]	193 [21]	36 [12-73]	8.39
MF	192	36.5 [16.5-59.5]	3420 [409]	15 [5-28]	7.92
+ OAH	144	48 [23-68]	2181 [284]	17 [8-37]	7.68
+ OAH + other	64	61.5 [33.5-75]	914 [114]	26 [14-49]	8.02
+ other (no OAH)	83	53 [26-68]	1158 [167]	26 [9-45]	6.93
+ other INCS	7	33 [0-77]	113 [21]	20 [6-79]	5.38
FP	33	44 [30-65]	156 [55]	23.5 [3.5-52]	2.83
+ OAH	34	56 [40-67]	305 [64]	19 [10-46]	4.77
+ OAH + other	14	52.5 [45-80]	60 [21]	54 [24.5-82.5]	2.89
+ other (no OAH)	13	41 [31-59]	121 [22]	22 [18-41]	5.50
+ other INCS	3	4 [0-65]	127 [11]	22 [8-48]	11.55

692 \*: monotherapy

693 FF: Fluticasone Furoate, FP: Fluticasone Propionate, MF: Mometasone Furoate, AZeFlu:  
 694 Azelastine-Fluticasone Propionate

695 p25: 25<sup>th</sup> percentile; p75: 75<sup>th</sup> percentile

696

697 **Table 3. Day 1 versus non-consecutive days**

	Day 1		1 <sup>st</sup> non-consecutive day		Other non-consecutive day		P value*
	N	VAS global, median [p25-p75]	N	VAS global, median [p25-p75]	N	VAS global, median [p25-p75]	Day 1 vs 1 <sup>st</sup> non-consecutive day
<b>All days</b>	4132	34 [12-60]	4132	25 [7-51]	24680	12 [2-32]	<0.001
<b>No treatment</b>	2214	26 [7-51]	2154	18 [4-44]	13651	8 [0-24]	<0.001
<b>AzeFlu</b>	162	44 [19-69]	187	26 [9-55]	1566	17 [6-35]	<0.001
<b>Other INCS treatment</b>	555	43 [22-64]	601	30 [11-55]	3403	17 [6-38]	<0.001

698 \*Statistical analysis by Wilcoxon and Mann-Whitney test

699 p25: 25<sup>th</sup> percentile; p75: 75<sup>th</sup> percentile

700

701 **Figure 1. Flow-chart of the study population**

702 **Figure 2: Percentage of days in each category of INCS treatment (first day and full data set)**

703 **Figure 3: Percentage of days in each category of treatment for VAS global measured (full**  
704 **dataset)**

705

706




Fig 1







Fig 2




Carrier 12:32

< Menu About us

The Allergy Diary was developed in collaboration between MACVIA-LR and ARIA.

**MACVIA\*LR**

MACVIA-LR (Contre les Maladies Chroniques pour un Vieillissement Actif en Languedoc-Roussillon, France) is a reference site of the European Innovation Partnership on Active and Healthy Ageing aimed at fighting chronic disease.




The ARIA (Allergic Rhinitis and its Impact on Asthma) initiative aims to educate and implement evidenced-based management of allergic rhinitis in conjunction with asthma

Carrier 12:34

< Menu My symptoms

Overall how much are your allergic symptoms bothering you today?

Not at all bothersome Extremely bothersome



Next

How much are your nose symptoms bothering you today?

How much are your eye symptoms bothering you today?

How much are your asthma symptoms bothering you today?

**23 countries (+3)**  
**16 languages**  
**22,500 users**  
**160,000 VAS days**  
**GPDR (May 25, 2018)**

Carrier 12:28

< Medication Nasal Done

QUICK SEARCH

Avamys (Fluticasone furoate)

Beclo aqua (Beclometasone)

Beclometasone (Beclometasone)

Beconase (Beclometasone)


I The I'm

Q W E R T Y U I O P


A S D F G H J K L

↑ Z X C V B N M ↵

123 ☺ space return



F3 online



F4 online




Fig 5A online

Eye symptoms




Fig 5B online

Asthma





Fig 5C online



None: No treatment  
FF: Fluticasone furoate  
OAH: Oral H1-anti-histamine  
AzeFlu: Azelastine-fluticasoen  
MF: Mometasone furoate



# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

**Table 1 online: Variations in VAS global levels within the same day**

Days with >1 VAS	N days	VAS global, median [p25-p75]		P value*
		1 <sup>st</sup> entry	2 <sup>nd</sup> entry	1 <sup>st</sup> vs 2 <sup>nd</sup> entry
<b>All days</b>	1,576	18 [4-45]	22 [6-50]	0.01
<b>Days without treatment</b>	866	14 [0-36]	17 [3-42]	0.005
<b>Days with AzeFlu treatment</b>	140	13 [4-41.5]	14 [4.5-53]	0.58
<b>Days with other INCS treatment</b>	177	29 [8-51]	25 [9-54]	0.90

\*Statistical analysis by Wilcoxon and Mann-Whitney test

p25: 25<sup>th</sup> percentile; p75: 75<sup>th</sup> percentile

**Figure 1 online: The Allergy Diary**

**Figure 2 online: Groups of users studied and excluded in the first analysis**

**Figure 3 online. Age distribution**

**Figure 4 online. VAS reporting trajectories in French users (n=520 users, 3,114 days)**

**Figures 5A, B, C online: Percentage of days in each category of treatment for VAS “eye”,  
“asthma” and “work” (full dataset)**